

Original Research Article

Risk factors of acute heart failure after the first episode of acute coronary syndrome

Sai Deepthi Janaki Rani Ivaturi^{1*}, Adithya Andanappa², Karthiga Vasudevan³,
Shashank Narayanappa⁴, Paidi Shrivatsam⁵

¹Gandhi Medical College, Hyderabad, Telangana, India

²Internal Medicine, Mysore Medical College and Research Institute, Mysore, Karnataka, India

³Yerevan State Medical University, Yerevan, Armenia

⁴Internal Medicine, Basaveshwara Medical College and Hospital, Chitradurga, Karnataka, India

⁵Government Medical College, Srikakulam, Andhra Pradesh, India

Received: 17 December 2024

Accepted: 24 February 2025

*Correspondence:

Dr. Sai Deepthi Janaki Rani Ivaturi,

E-mail: deepthiivaturi2699@gmail.com

Copyright: © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

Background: Factors that speed up the development of acute heart failure (AHF) after acute coronary syndrome (ACS), are understudied in developing country like India. Therefore, this study was aimed to determine the risk factor of heart failure (HF) among the post-ACS patients.

Methods: This descriptive cross-sectional study was performed at Government General Hospital (GGH) Srikakulam, Andhra Pradesh, India. 108 patients who were admitted with HF cardiology unit of GGH after ACS from September 2023 to November 2024 were enrolled in the study. An established inclusion and exclusion criteria along with consecutive sampling technique were used for patients' recruitment. A self-structured proforma was applied to gather data. Data analysis was done in the statistical package for the social sciences (SPSS) version 25.0.

Results: HF was more prevalent among those post-acute coronary syndrome patients who had an age group with 40 years or above (69.23%), male gender (62.82%), anterior wall myocardial infarction (38.46%), reduced ejection fraction (67.90%), left anterior descending artery pathology (17.95%), absence of myocardial revascularization procedure (46.15%), diabetes mellitus (52.56%), hypertension (78.20%), anemia (53.85%), hyperlipidaemia (75.64%), and who had history of smoking (51.28%). A significant number of patients also had chronic kidney disease (23.10%).

Conclusions: Factors such as an age group with 40 years or above, male gender, anterior wall myocardial infarction, reduced ejection fraction, left anterior descending artery pathology, absence of myocardial revascularization procedure, diabetes mellitus, hypertension, anemia, hyperlipidemia and history of smoking, all accelerate the development of HF among post-ACS patients.

Keywords: Acute, Coronary, Factors, Failure, Heart, Risk, Syndrome

INTRODUCTION

Acute coronary syndrome (ACS) refers to a spectrum of clinical presentations resulting from a sudden reduction in myocardial perfusion, which manifests as symptoms such as severe chest pain radiating to the jaw or left arm, tachycardia, diaphoresis, dyspnea, dizziness and gastrointestinal discomfort, including nausea or

vomiting.^{1,2} The primary pathological event in ACS involves the rupture or erosion of an unstable atherosclerotic plaque, leading to partial or complete thrombotic occlusion of a coronary artery. The condition is classified into three subtypes based on electrocardiographic findings, unstable angina (UA) and non-ST-segment elevation myocardial infarction (NSTEMI), which lack ST-segment elevation and acute

ST-segment elevation myocardial infarction (STEMI), characterized by the hallmark ST-segment elevation on ECG.^{3,4} ACS is a life-threatening condition with a rapid onset, accelerated progression and poor prognosis. Its clinical course is often complicated by a variety of severe outcomes, including heart failure (HF), which is among the most prevalent complications.^{5,6} The pathogenesis of HF post-myocardial infarction involves a cascade of events, including myocardial necrosis, myocardial stunning and mechanical complications such as papillary muscle rupture, ventricular septal defect and ventricular free wall rupture. Cellular damage begins within minutes of ischemia, with structural alterations and edema evident by 30 minutes and irreversible myocyte death occurring within 3 hours. Additionally, reperfusion therapy, while essential, contributes to injury through the generation of reactive oxygen species, exacerbating myocyte loss. The resulting inflammatory cascade further promotes HF development.^{7,8}

The presence of comorbidities such as diabetes mellitus, hypertension, anemia, chronic kidney disease, Hyperlipidemia and smoking exacerbates the risk and progression of HF following ACS.⁹⁻¹⁴ Early identification and management of these risk factors could play a critical role in mitigating HF development and improving clinical outcomes in patients recovering from ACS. This study, therefore, seeks to investigate the potential risk factors contributing to HF onset in patients post-ACS, with the goal of reducing morbidity and improving prognostic outcomes.

METHODS

Study design

This descriptive cross-sectional study.

Study place

was performed in the Cardiology Unit of Government general hospital (GGH) Srikakulam Andhra Pradesh India.

Study duration

from September 2023 to November 2024 were enrolled in the study.

Study population

108 patients who were admitted with AHF cardiology unit of GGH after ACS. An established inclusion and exclusion criteria along with consecutive sampling technique were used for patients' recruitment.

Inclusion criteria

All patients were included in the study who had, age 18 years or above, admitted in cardiac unit of GGH with

diagnosed HF following ACS, history of only one episode of diagnosed ACS in last one year with complete medical record of their first admission with ACS.

Exclusion criteria

While those admitted patients who had, history of stable angina, congenital heart disease, severe liver disease, autoimmune disease, active infectious disease, oncological conditions and who were not willing to participate were excluded from the study.

ACF and acute coronary syndrome

To determine whether patients had AHF, the diagnostic criteria for AHF were based on the European Society of Cardiology's 2021 guidelines for the diagnosis and treatment of acute and chronic heart failure. AHF was diagnosed based on clinical symptoms such as dyspnea, fast heartbeat, third heart sound, elevated jugular vein pressure, lung rales and lower limb edema. Furthermore, higher B-type natriuretic peptide (BNP) levels were taken into account. Similarly, ACS was diagnosed using the European society of cardiology's 2023 guidelines for the diagnosis and management of ACS. The diagnosis and classification of ACS was made on the basis of clinical features (chest pain (crushing, pressure like feeling in character) with radiation to jaw or left arm, racing heartbeat, shortness of breath, sudden sweating, dizziness and nausea or vomiting), electrocardiographic (ECG), cardiac biomarker (enzymes) levels, coronary angiography and echocardiography were also taken into account.

Data collection

A self-designed questionnaire was applied for data collection. It had three components. First component had sociodemographic features such as age (less than 40 years and 40 and above years), gender (male or female) and comorbidities presence or absence (diabetes mellitus, hypertension, anemia, chronic kidney disease, hyperlipidemia and smoking).

Second component was regarding the variables that were associated with findings of ECG (location of MI), coronary angiography (vessel involved) and echocardiography (ejection fraction: normal or reduced) during ACS before HF development. Third component was comprised of biochemical markers' values such as serum lipid level, haemoglobin, serum creatinine level and cardiac biomarkers. Information about the revascularization (thrombolysis, percutaneous coronary intervention, coronary artery bypass, None) was also noted in this part of the questionnaire.

Statistical analysis

The statistical analysis of data was performed through statistical package for the social sciences (SPSS) 25.0

(Armonk, NY: IBM Corp.). By using descriptive statistics, numerical data were shown as mean±standard deviation and nominal data were presented as frequencies and percentages.

RESULTS

Means of various study variables such as age and ejection fraction were 59.09 with a standard deviation (SD) of±12.12 year and 42.58 with SD of±8.10 respectively. Table 1 indicates that that incidence of HF was more common among the patients who had older age group (age 40 years or above) and male gender in contrast to the patients who had younger age group (age less than 40 years) and female gender. Table 2 shows that frequency of HF was higher among the participants who had ECG changes in anterior territory followed by inferior, lateral, septal, anterior and inferior, anterior and lateral and inferior and lateral territories. It also displays that patients with reduced ejection fraction, single-vessel disease which involved mainly left anterior descending artery, two-vessel disease involved left anterior descending artery and circumflex coronary artery and with none myocardial revascularization procedures, had higher HF frequency in comparison to patients with normal ejection

fraction, single-vessel disease which involved either right coronary artery or circumflex artery, two vessel disease involved either left anterior descending artery and right coronary artery or circumflex coronary artery and right coronary artery and with different myocardial revascularization procedures, had lower HF frequency.

Table 1: Demographic characteristics of the study population.

Variables	Frequency (N)	%
Age (in years)		
Less than 40	34	30.77
40 and above	74	69.23
Gender		
Male	68	62.82
Female	40	37.18

Table 3 manifests that HF incidence was more prevalent among the patients with who had diabetes mellitus, hypertension, anemia, hyperlipidemia and history of smoking as compared to who had not above-mentioned comorbidities. However, a significant number of patients also had chronic kidney disease.

Table 2: Frequency and percentages of cardiac risk factors in study population.

Variables	Frequency (N)	%
ECG territory changes		
Anterior	41	38.46
Inferior	26	24.36
Lateral	12	11.54
Septal	11	10.25
Anterior and Inferior	8	7.70
Anterior and lateral	5	5.13
Inferior and lateral		2.56
Ejection fraction on echocardiography		
Normal	35	32.10
Reduced	73	67.90
Coronary angiography		
Not performed	51	47.44
Single-vessel disease	33	30.77
Left descending artery	19	17.95
Right coronary artery	8	7.70
Circumflex artery	5	5.13
Two-vessel disease	17	16.70
Left anterior descending artery and circumflex coronary artery	10	8.97
Left anterior descending artery and right coronary artery	5	5.13
Circumflex coronary artery and right coronary artery	3	2.56
Triple-vessel disease	5	5.13
Myocardial revascularization		
Thrombolysis	20	17.95
Percutaneous coronary intervention	31	29.50
Coronary artery bypass graft	7	6.41
Not performed	50	46.15

Table 3: Frequency and percentages of systemic risk factors in study population.

Variables	Frequency (N)	%
Diabetes mellitus		
Yes	56	52.56
No	50	47.44
Hypertension		
Yes	84	78.20
No	22	21.80
Anemia		
Yes	57	53.85
No	50	46.15
Chronic kidney disease		
Yes	24	23.10
No	83	76.90
Hyperlipidemia		
Yes	81	75.64
No	26	24.36
History of smoking		
Yes	55	51.28
No	51	48.72

DISCUSSION

ACS and HF

ACS is a life-threatening condition characterized by sudden onset, rapid progression and a grim prognosis. It is associated with severe complications, among which HF is one of the most prevalent and deleterious outcomes. This study offers significant insights into the risk factors contributing to HF development following ACS.

Heart failure commonly arises after myocardial infarction (MI) hospitalization due to cardiomyocyte death and subsequent scar tissue formation. This process initiates chronic neurohumoral activation involving the renin-angiotensin-aldosterone system and sympathetic nervous system upregulation. These mechanisms promote ventricular remodelling, altering ventricular geometry and resulting in wall thinning, ischemic mitral regurgitation and additional cardiomyocyte loss. Such pathological changes underline the progression of HF after ACS.^{7,12,14}

Sociodemographic factors influencing HF after ACS

The study analyzed how sociodemographic factors in the research population impacted HF development post-ACS. It was observed that older age groups had a higher incidence of HF following ACS. These findings are consistent with a similar study conducted in the United Kingdom, which highlighted age as a significant determinant of HF progression.⁴ Regarding gender differences, this study found a higher HF rate among male patients, corroborating findings from a study conducted in Romania.¹ However, contrasting evidence from another study reported a predominance of HF

among females after ACS.⁵ These discrepancies may stem from variations in racial, sociodemographic and geographical characteristics across different populations.

Cardiac risk factors for HF post-ACS

The location and extent of ACS significantly influenced the development of HF. Patients with ACS involving the anterior wall of the heart and the left anterior descending artery exhibited a higher HF frequency compared to ACS affecting other regions. This can be attributed to the larger extent of irreversible left ventricular damage associated with anterior wall ACS due to severe lesions in the left anterior descending artery. Several studies align with this finding, affirming the higher risk linked to this specific anatomical location.^{6,9}

Additionally, a reduced ejection fraction was strongly associated with an increased likelihood of HF development, a conclusion supported by numerous global studies.^{1,3,9} Furthermore, patients who did not undergo myocardial revascularization procedures demonstrated a higher rate of post-ACS HF, consistent with findings from various international studies.^{4,5}

Systemic risk factors of HF following ACS

The study identified several systemic risk factors contributing to HF development post-ACS. The presence of comorbidities such as diabetes mellitus, hypertension, anemia, hyperlipidemia and smoking history significantly increased HF rates compared to patients without these comorbidities. These observations are well-supported by literature emphasizing the role of comorbidities in accelerating HF onset.⁸⁻¹³ Comorbidities exacerbate HF development through mechanisms like accelerated

vascular atherosclerosis, microvascular injury and impaired left ventricular remodelling, all of which contribute to poor cardiac outcomes. Such mechanisms underscore the importance of managing systemic risk factors in patients with ACS to mitigate HF risk.

This study's primary strength lies in its originality, as it is the only research in the study's locality to identify potential risk factors for HF in patients experiencing one episode of ACS. This localized focus provides valuable insights for regional healthcare planning and targeted interventions.

However, the study is not without limitations. Being a single-center study with a relatively small sample size, its findings may be subject to bias and limited generalizability. Future research involving multi-center studies with larger, more diverse populations is encouraged to validate these findings and enhance their applicability.

CONCLUSION

This present study results have suggested that frequency of heart failure in the post-acute coronary syndrome patients was higher among those who had the age 40 years or above and male gender. The main risk factors associated with the development of HF were anterior wall myocardial infarction, left anterior descending artery pathology, reduced ejection fraction, absence of myocardial revascularization procedure and presence of comorbidities such as diabetes mellitus, hypertension, anemia, hyperlipidemia and history of smoking.

A significant number of patients also had chronic kidney disease. Health officials should raise people's awareness of risk factors for heart failure following acute coronary syndrome through public service announcements, regular health fairs, newsletters and mass media campaigns. Heart failure incidence could be reduced by increasing public knowledge about risk factors and preventive actions. Furthermore, a proper assessment and forecasting could limit the risk of HF in post-ACS patients. It could also effectively overcome post-ACS problems and maximize their degree of recovery.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

1. Bustea C, Tit DM, Bungau AF, Bungau SG, Pantea VA, Babes EE, et al. Predictors of readmission after the first acute coronary syndrome and the risk of recurrent cardiovascular events—seven years of patient follow-up. *Life*. 2023;13(4):950.
2. Keykhaei M, Ashraf H, Rashedi S, Farrokhpour H, Heidari B, Zokaei S, et al. Differences in the 2020 ESC versus 2015 ESC and 2014 ACC/AHA guidelines on the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Curr Atherosclerosis Rep*. 2021;23:1-10.
3. Zhang L, Hailati J, Ma X, Liu J, Liu Z, Yang Y, et al. Analysis of risk factors for different subtypes of acute coronary syndrome. *J Int Med Res*. 2021;7:67-9.
4. Cahill TJ, Kharbanda RK. Heart failure after myocardial infarction in the era of primary percutaneous coronary intervention: Mechanisms, incidence and identification of patients at risk. *World J Cardiol*. 2017;9:407-5.
5. Arora S, Stouffer GA, Kucharska-Newton AM, Qamar A, Vaduganathan M, Pandey A, et al. Twenty year trends and sex differences in young adults hospitalized with acute myocardial infarction: the ARIC Community Surveillance Study. *Circulation*. 2019;139(8):1047-56.
6. Liang J, Zhang Z. Predictors of in-hospital heart failure in patients with acute anterior wall ST-segment elevation myocardial infarction. *Int J Cardiol*. 2023;375:104-9.
7. Harjola VP, Parissis J, Bauersachs J, Brunner-La Rocca HP, Bueno H, Čelutkienė J, et al. Acute coronary syndromes and acute heart failure: a diagnostic dilemma and high-risk combination. A statement from the Acute Heart Failure Committee of the Heart Failure Association of the European Society of Cardiology. *Eur J Heart Failure*. 2020;22(8):1298-314.
8. Ulvenstam A, Graipe A, Irewall AL, Söderström L, Moee T. Incidence and predictors of cardiovascular outcomes after acute coronary syndrome in a population-based cohort study. *Sci Rep*. 2023;13(1):3447.
9. Jenča D, Melenovský V, Stehlik J, Staněk V, Kettner J, Kautzner J, et al. Heart failure after myocardial infarction: incidence and predictors. *ESC Heart Failure*. 2021;8(1):222-37.
10. Okkonen M, Havulinna AS, Ukkola O, Huikuri H, Pietilä A, Koukkunen H, et al. Risk factors for major adverse cardiovascular events after the first acute coronary syndrome. *Ann Med*. 2021;53(1):817-23.
11. Gouda P, Savu A, Baaney KR, Kaul P, Welsh RC. Long-term risk of death and recurrent cardiovascular events following acute coronary syndromes. *PLoS One*. 2021;16(7):254008.
12. Baluja A, Rodríguez-Mañero M, Cordero A, Kreidieh B, Iglesias-Alvarez D, García-Acuña JM, et al. Prediction of major adverse cardiac, cerebrovascular events in patients with diabetes after acute coronary syndrome. *Diabetes Vasc Dis Res*. 2020;17(1):1479164119892137.
13. Chia JE, Ang SP. Drinking patterns of alcohol and risk of major adverse cardiovascular events after an acute coronary syndrome. *Eur J Prev Cardiol*. 2024;19:780.

14. Zhang F, Mohamed MO, Ensor J, Peat G, Mamas MA. Temporal trends in comorbidity burden and impact on prognosis in patients with acute coronary syndrome using the Elixhauser Comorbidity Index Score. *Am J Cardiol*. 2020;125(11):1603-11.
15. Ahmad Y, Petrie MC, Jolicœur EM, Madhavan MV, Velazquez EJ, et al. PCI in patients with heart failure: current evidence, impact of complete revascularization, and contemporary techniques to

improve outcomes. *J Soc Cardiovasc Angiogr Interv*. 2022;1(2):100020.

Cite this article as: Ivaturi SDJR, Andanappa A, Vasudevan K, Narayanappa S, Shrivatsam P. Risk factors of acute heart failure after the first episode of acute coronary syndrome. *Int J Clin Trials* 2025;12(2):105-10.